

Hausdorff dimension and anyonic distribution functions

Wellington da Cruz*

Departamento de Física,

Universidade Estadual de Londrina, Caixa Postal 6001,

Cep 86051-970 Londrina, PR, Brazil

(February 1, 2008)

Abstract

We obtain the distribution functions for anyonic excitations classified into equivalence classes labeled by Hausdorff dimension, h and as an example of such anyonic systems, we consider the collective excitations of the Fractional Quantum Hall Effect (FQHE).

PACS numbers: 05.30.-d, 05.70Ge

Keywords: Distribution functions; Hausdorff dimension; Anyonic excitations; Fractional quantum Hall effect

Typeset using REVTeX

*E-mail: wdacruz@fisica.uel.br

We have obtained in [1] from a continuous family of Lagrangians for fractional spin particles a path integral representation for the propagator and its representation in moment space. On the other hand, the trajectories swept out by scalar and spinning particles can be characterized by the fractal parameter h (or the Hausdorff dimension). We have that, L , the length of closed trajectory with size R has its fractal properties described by $L \sim R^h$ [2], such that for scalar particle $L \sim \frac{1}{p^2}$, $R^2 \sim L$ and $h = 2$; for spinning particle, $L \sim \frac{1}{p}$, $R^1 \sim L$ and $h = 1$. From the form of anyonic propagator given in [1], with the spin defined into the interval $0 \leq s \leq 0.5$, we have extracted the following formula, $h = 2 - 2s$, which relates the Hausdorff dimension h and the spin s of the particle. Thus, for anyonic particle, h takes values within the interval $1 < h < 2$. In [3], we have classified the fractional spin particles or anyonic excitations in terms of equivalence classes labeled by h . Therefore, such particles in a specific class can be considered on equal footing.

On the other hand, in the context of FQHE systems, the filling factor, a parameter which characterize that phenomenon, can be classified in this terms too. We have, therefore, a new hierarchy scheme for the filling factors [4], which is extracted from the relation between h and the statistics ν (or the filling factors). Our approach, contrary to literature [5] do not have an empirical character and we can predicting for which values of ν FQHE can be observed. A Braid group structure behind this classification also was noted [6] and in [3] a topological invariant, $\mathcal{W} = h + 2s - 2p$, was introduced which relates a characteristic of the multiply connected spaces with the numbers of fractional spin particles, p , and the quantities related to the particles, h and s (note that this invariant has a connection with the *Euler characteristic* of the surface). The anyonic model, a charge-flux system, constitutes a topological obstruction (holes with hard core repulsion) in this description and the elements of the Braid group are equivalent trajectories.

Now, we propose a statistical weight for such excitations in terms of h , as follows:

$$\omega_j = \frac{[G_j + (N_j - 1)(h - 1)]!}{N_j! [G_j + (N_j - 1)(h - 1) - N_j]!}, \quad (1)$$

where G_j means a group of quantum states, N_j is the number of particles and for, $h = 2$ we have bosons and for $h = 1$ we have fermions. For, $1 < h < 2$, we have anyonic excitations which interpolates between these two extremes. The fractal parameter h_i (i means a specific interval) is related to statistics ν , in the following way:

$$\begin{aligned} h_1 &= 2 - \nu, & 0 < \nu < 1; & & h_2 &= \nu, & 1 < \nu < 2; \\ h_3 &= 4 - \nu, & 2 < \nu < 3; & & h_4 &= \nu - 2, & 3 < \nu < 4; \\ h_5 &= 6 - \nu, & 4 < \nu < 5; & & h_6 &= \nu - 4, & 5 < \nu < 6; \\ h_7 &= 8 - \nu, & 6 < \nu < 7; & & h_8 &= \nu - 6, & 7 < \nu < 8; \\ h_9 &= 10 - \nu, & 8 < \nu < 9; & & h_{10} &= \nu - 8, & 9 < \nu < 10; \\ &etc, \end{aligned} \quad (2)$$

such that this spectrum of ν , as we can see, has a complete mirror symmetry and for a given h , we collect different values of ν in a specific class, for example, consider $h = \frac{5}{3}$ and $h = \frac{4}{3}$, so we obtain; $\left\{ \frac{1}{3}, \frac{5}{3}, \frac{7}{3}, \frac{11}{3}, \frac{13}{3}, \frac{17}{3}, \dots \right\}_{h=\frac{5}{3}}$ and $\left\{ \frac{2}{3}, \frac{4}{3}, \frac{8}{3}, \frac{10}{3}, \frac{14}{3}, \frac{16}{3}, \dots \right\}_{h=\frac{4}{3}}$.

These particles or excitations, in the class, share some common characteristics, according our interpretation, in the same way that bosons and fermions. We observe that each interval

contributes with one and only one particle to the class and the statistics and the spin, are related by $\nu = 2s$.

We stress that, our expression for statistical weight Eq.(1), is more general than an expression given in [7], once if we consider the relation between h and ν , we just obtain that first result and extend it for the complete spectrum of statistics ν , for example:

$$\begin{aligned}\omega_j &= \frac{[G_j + (N_j - 1)(1 - \nu)]!}{N_j! [G_j + (N_j - 1)(1 - \nu) - N_j]!}, & 0 < \nu < 1; \\ \omega_j &= \frac{[G_j + (N_j - 1)(\nu - 1)]!}{N_j! [G_j + (N_j - 1)(\nu - 1) - N_j]!}, & 1 < \nu < 2; \\ \omega_j &= \frac{[G_j + (N_j - 1)(3 - \nu)]!}{N_j! [G_j + (N_j - 1)(3 - \nu) - N_j]!}, & 2 < \nu < 3; \\ \omega_j &= \frac{[G_j + (N_j - 1)(\nu - 3)]!}{N_j! [G_j + (N_j - 1)(\nu - 3) - N_j]!}, & 3 < \nu < 4; \\ &etc.\end{aligned}\tag{3}$$

The expressions Eq.(3) were possible because of the mirror symmetry as just have said above. But, our approach in terms of Hausdorff dimension, h , have an advantage, because we collect into equivalence class the anyonic excitations and so, we consider on equal footing the excitations in the class and this is a new approach which enter on considerations about fractional spin particles. For different species of particles, the statistical weight take the form

$$\Gamma = \prod_j \omega_j = \prod_j \frac{[G_j + (N_j - 1)(h - 1)]!}{N_j! [G_j + (N_j - 1)(h - 1) - N_j]!},\tag{4}$$

which reduces to Eq.(1) for only one species. Now, we can consider the entropy for a given class h . Taking the logarithm of Eq.(4), with the condition that, N_j and G_j are vary large numbers and defining the average occupation numbers, $n_j = \frac{N_j}{G_j}$, we obtain for a gas not in equilibrium, an expression for the entropy

$$\begin{aligned}\mathcal{S} &= \sum_j G_j \left\{ [1 + n_j(h - 1)] \ln \left(\frac{1 + n_j(h - 1)}{1 + n_j(h - 1) - n_j} \right) \right. \\ &\quad \left. - n_j \ln \left(\frac{n_j}{1 + n_j(h - 1) - n_j} \right) \right\},\end{aligned}\tag{5}$$

such that for $h = 2$ and $h = 1$, we obtain the expressions for a Bose and Fermi gases not in equilibrium [8],

$$\mathcal{S} = \sum_j G_j \{ (1 + n_j) \ln (1 + n_j) - n_j \ln n_j \};\tag{6}$$

$$\mathcal{S} = - \sum_j G_j \{ n_j \ln n_j + (1 - n_j) \ln (1 - n_j) \},\tag{7}$$

respectively.

The distribution function for a gas of the class h can be obtained from the condition of the entropy be a maximum. Thus, we have

$$n_j \xi = \{1 + n_j(h - 1)\}^{h-1} \{1 + n_j(h - 2)\}^{2-h}, \quad (8)$$

where $\xi = \exp \{(\epsilon_j - \mu)/KT\}$, has the usual definition and the Bose and Fermi distributions are obtained for values of $h = 2, 1$; respectively.

In the context of the Fractional Quantum Hall Effect (FQHE), as we said in the introduction, the filling factor (rational number with an odd denominator), can be also classified in terms of h [4]. We have that the anyonic excitations are collective excitations manifested as quasiparticles or quasiholes in FQHE systems. Thus, for example, we have the collective excitations as given in the beginning for $h = \frac{5}{3}$ and $h = \frac{4}{3}$. Now, we have noted that these collections contain filling fractions, in particular, $\nu = \frac{1}{3}$ and $\nu = \frac{2}{3}$, that experimentally were observed [9] and so we are able to estimate for which values of ν the largest charge gaps occurs, or alternatively, we can *predicting* FQHE. As was observed in [4] this is a new schem hierarchy scheme for the filling factors, that express the occurrence of the FQHE in more fundamental terms, that is, relating the fractal parameter h and the filling factors ν . Therefore, for anyonic excitations in a stronger magnetic field at low temperatures, our generalization (Eq.3) of the results in [7] can be considered.

In summary, we have obtained distribution functions (Eq.8) for anyonic excitations in terms of the Hausdorff dimension h , which classified the anyonic excitations into equivalence classes and reduces to fermionic and bosonic distributions, when $h = 1$ and $h = 2$, respectively. In this way, we have extended (Eq.3) results of the literature [7] for the complete spectrum of statistics ν . A connection with the FQHE, considering the filling factors into equivalence classes labeled by h also was considered and an estimate for occurrence of FQHE made.

REFERENCES

- [1] W. da Cruz, preprint/UEL-DF/W-01/97, hep-th/9803020.
- [2] A. M. Polyakov, in *Proc. Les Houches Summer School* **vol. II**, ed. E. Brézin and J. Zinn-Justin (North Holland, 1990), 305.
- [3] W. da Cruz, preprint/UEL-DF/W-02/97, hep-th/9803043; *ibid*, preprint/UEL-DF/W-03/97, hep-th/9802135.
- [4] W. da Cruz, preprint/UEL-DF/W-04/97, cond-mat/9802266.
- [5] F. D. M. Haldane, Phys. Rev. Lett. **51** (1983) 605; B. I. Halperin, Phys. Rev. Lett. **52** (1984) 1583; see also [9].
- [6] W. da Cruz, preprint/UEL-DF/W-05/97, cond-mat/9802267.
- [7] Y. S. Wu, Phys. Rev. Lett. **73** (1994) 922 .
- [8] E. M. Lifshitz and L. P. Pitaevskii, *Statistical Physics, Part 1, 3rd. edition* (Pergamon Press, Oxford, 1980).
- [9] A. H. MacDonald, cond-mat/9410047; C. Cros and A. H. MacDonald, Phys. Rev. **B 42** (1990) 10811 and references therein.